## Copyright (C) 2011 David K. Levine

This document is an open textbook; you can redistribute it and/or modify it under the terms of version 1 of the open text license amendment to version 2 of the GNU General Public License. The open text license amendment is published by Michele Boldrin et al at http://levine.sscnet.ucla.edu/general/gpl.htm; the GPL is published by the Free Software Foundation at <a href="http://www.gnu.org/copyleft/gpl.html">http://www.gnu.org/copyleft/gpl.html</a>.

If you prefer you may use the Creative Commons attribution license http://creativecommons.org/licenses/by/2.0/

# The Evolution and Cooperation

# The Standard Setting

- playing repeatedly against a fixed opponent vs. myopia
- pick a players at random from a large population everyone sees play
- players randomly matched, results of all matches revealed anonymously
- players matched randomly see results only of own match (this is how experiments are conducted)
- no issue of causality

# **Learning versus Evolution**

- · learning: beliefs updated based on experience
- evolution: better strategies do better/random mutation, no beliefs

# **Dynamics: Best Response Dynamics**

- discrete time best response
- discrete time partial best response
   -individual vs population model
- continuous time best response

2,2	0,0
0,0	1,1

mixed equilibrium 1/3-2/3

illustrate three dynamics

# Shapley example

0,0	1,2	2,1
2,1	0,0	1,2
1,2	2,1	0,0

note that (0,0) is never hit, but always in Nash equilibrium

"smoothed best response" saddles and medium run

# **Dynamics: Replicator**

- definition
- as a model of social learning
- as a stimulus-response model
- probability matching issues

# Kandori-Mailath-Rob Young and the Ultra Long-Run

1>x>y>0

X,X	y,0
0,y	1,1

p=pr(x)=(1-y)/(x-y+1) (indifference between up and down)

1 is pareto efficient

x is risk dominant if and only if 1<x+y

for example, x=3/4, y=1/2

# finite population of N players

- deterministic dynamic
- mutations
- 1-p mutations x->1
- p mutations x->1
- relative waiting times

#### Comments:

Nachbar: it can take a long time to learn to eliminate dominated strategies (deterministic dynamic)

Ellison: the very long run can be very long, but much shorter with local interaction

Johnson, Pesendorfer and Levine

# **Overview**

- The problem of multiple equilibria and the folk theorem
- One answer: an evolutionary approach to see what equilibrium results

- Propogation through imitation rather than innovation
- Strategies that depend on the opponent's type in a random matching game
- Example: cooperate if opponent is same type, punish if he is different

   usual folk theorem result, this is an equilibrium, but so is never
   cooperate
- No errors: long-run equilibrium is efficient, but bursts of conflict in which players minmax the difference in their payoffs
- Errors in an additively separable environment: maximize a weighted sum of own and opponents payoff – preferences for altruism and spite

#### Literature

- Kandori, Mailath and Rob [1993] and Young [1993]
- Morris, Rob and Shin [1993] one-half dominance
- Ellison [2000] co-radius
- Bergin and Lipman [1994] relative probabilities of noise matters
- Johnson, Pesendorfer and Levine [2000] emergence of cooperation in a trading game
- Kandori and Rob [1993] winning pairwise contests sufficient because implies 1/2-dominance

# The Model and Basic Characterization of Long Run Outcomes

- symmetric normal form game
- single population of players
- finitely many pure strategies  $s \in S$
- mixed strategies  $\sigma \in \Sigma$
- $\bullet$  utility on his own pure strategy and mixed strategy of population  $u(s,\sigma)$
- $u(s,\sigma)$  continuous in  $\sigma$
- two player matching games:  $u(s,\sigma)$  linear in  $\sigma$

## **Evolution**

- m player population
- each plays a pure strategy
- $\bullet$  distribution of strategies at t denoted by  $\sigma_t \in \Sigma$
- $\bullet$  initial distribution  $\sigma_0$

#### The Imitative Process

- $\sigma_t$  is determined from  $\sigma_{t-1}$  according to the "imitative" process
- 1) One player i chosen at random; only this player changes strategy
- 2) probability  $C\varepsilon$  imitation strategies chosen in proportion to previous period frequency player i chooses from S randomly using the probabilities  $\sigma_{t-1}$ .
- 3) probability  $\varepsilon^n$  innovation strategies are entirely at random -player i chooses each strategy from S with equal probability
- 4) probability  $1 C\varepsilon \varepsilon^n$  relative best response best response among those strategies that are actually used player i randomizes with equal probability among the strategies that solve

$$\max_{s \in \text{supp}(\sigma_{t-1})} u(s, \sigma_{t-1})$$

#### The Markov Process

- The imitative process is a Markov process M on state space  $\Sigma^m \subset \Sigma$  of all mixed strategies consistent with the grid induced by each player playing a pure strategy
- process M is positively recurrent because of innovation
- ullet so M has unique invariant distribution  $\mu^{arepsilon}$
- goal: characterize  $\mu \equiv \lim_{\varepsilon \to 0} \mu^{\varepsilon}$

Assumption: imitation is much more likely than innovation.

Unlikely Innovation: n > m

as  $\varepsilon \to 0$  probability of *every* player changing strategy by imitation much greater than probability a single player innovates

## **Basic Results**

## Pure Strategies

- mixed strategies less stable than pure strategies
- mixed strategy can evolve to pure only using imitation
- pure cannot evolve at all without at least one innovation

**Theorem 1:**  $\mu = \lim_{\varepsilon} \mu = \lim_{\varepsilon} \mu$ 

#### Pairwise Contests

what it means to win pairwise contests:

 $0 \le \alpha \le 1$  mixed strategy that plays s with probability  $\alpha$  and  $\tilde{s}$  with probability  $1-\alpha$  denoted by  $\alpha s + (1-\alpha)\tilde{s}$ .

**Definition 1:** The strategy s beats  $\tilde{s}$  iff

$$u(s, \alpha s + (1 - \alpha)\tilde{s}) - u(\tilde{s}, \alpha s + (1 - \alpha)\tilde{s}) > 0$$

for all  $1/2 \le \alpha < 1$ 

**Definition 2:** The strategy s weakly beats  $\tilde{s}$  iff

$$u(s, \alpha s + (1 - \alpha)\tilde{s}) - u(\tilde{s}, \alpha s + (1 - \alpha)\tilde{s}) > 0$$

for all 
$$1/2 < \alpha < 1$$
 and  $u(s, \frac{1}{2}s + \frac{1}{2}\tilde{s}) - u(\tilde{s}, \frac{1}{2}s + \frac{1}{2}\tilde{s}) = 0$ 

**Definition 2':** The strategy s is **tied** with  $\tilde{s}$  iff

$$u(s, \alpha s + (1 - \alpha)\tilde{s}) - u(\tilde{s}, \alpha s + (1 - \alpha)\tilde{s}) = 0$$

*for all*  $1 \ge \alpha \ge 0$ 

**Definition 3**: If sbeats all  $\tilde{s} \neq s$  we say that s beats the field. If for all  $\tilde{s} \neq s$  either s weakly beats  $\tilde{s}$  or is tied with  $\tilde{s}$  we say that s weakly beats the field

## A Sufficient Condition for Long Run Equilibrium

**Theorem 2:** If m is sufficiently large and s beats the field then  $\mu(s) = 1$ . If m is sufficiently large and s weakly beats the field then  $\mu(s) > 0$ . Moreover, if  $\mu(\tilde{s}) > 0$  then  $u(s, \frac{1}{2}s + \frac{1}{2}\tilde{s}) - u(\tilde{s}, \frac{1}{2}s + \frac{1}{2}\tilde{s}) = 0$ 

Uses standard Kandori-Mailath-Rob-Young tree-trimming arguments

This condition is similar to ½ dominance, but much weaker – it isn't necessary to beat combinations of other strategies, just one at a time

We'll see how that gets used later

# Matching Games with Behavioral Types

- each period players matched into pairs to play a symmetric normal form game
- prior choosing an action, each player receives a "signal" about how his opponent will behave in the game
- how does the long-run outcome depends upon the amount of information contained in the signals?

## **Formalities**

- action space A
- $\bullet$  payoff of a player who takes action a and whose opponent takes action a ' is U(a,a ')
- ullet actual strategy space is a finite abstract space S

# strategies serve two roles

- 1. influence the information that is generated about the player and his opponent
- 2. govern the behavior as a function of the generated information

## Informational Function of Strategies

- each player receives a signal  $y \in Y$ , a finite set
- probability of signal given by  $\pi(y\mid s,s')$  if the player uses s and the opponent s'
- signals are private information and reflects what the opponent can learn about his opponent prior to the interaction

# Behavioral Function of Strategies

- each strategy s gives rise to a map  $s: Y \to A$
- several strategies may induce the same map, yet differ in the probability with which they send signals

for every map from signals to actions there is some strategy that induces that map

**Assumption 0:** If  $f: Y \to A$  there is a strategy  $s \in S$  with s(y) = f(y). the space of signals is necessarily smaller than the set of strategies the cardinality of the space of strategies is at least  $A^Y$ , which is greater than that of Y provided that there are at least two signals

#### **Motivation**

- People have some ability to detect lying
- Frank [1987] some people blush whenever they tell a lie
- A good lie is costly to construct Ben Gurion airport
- The signal is a combination of a self-report of intentions together with an involuntary signal about lying
- "I'll still respect you in the morning [blush]"

## example

$$Y = \{0,1\}$$
 
$$\pi(y = 0 \mid s',s) = 1 \text{ if } s' = s$$
 
$$\pi(y = 1 \mid s',s) = 1 \text{ if } s' \neq s$$

two players meet who use the same strategy then both receive the signal 0 whereas when two players meet who use different strategies then both receive the signal 1

players recognize if the opponent uses the same strategy or a different strategy prior to play

important, because it turns out that strategies that recognize themselves are likely to emerge in the long-run equilibrium.

signals affect payoffs only indirectly by affecting behavior player i uses strategy s and opponent uses strategy s' expected payoff of player i

$$\sum_{y' \in Y} \sum_{y \in Y} U(s(y), s'(y')) \pi(y \mid s, s') \pi(y' \mid s', s) \sigma(s')$$

the function  $u(s,\sigma)$ 

$$u(s,\sigma) = \sum_{s' \in S} \sum_{y' \in Y} \sum_{y \in Y} U(s(y), s'(y')) \pi(y \mid s, s') \pi(y' \mid s', s) \sigma(s')$$

## Perfect Identification

each player can identify with certainty whether the opponent is using the same strategy

assume for notational simplicity one signal, denoted by  $y_0$  that perfectly identifies that the opponent is using the same strategy.

**Assumption 1:** There is a  $y_0 \in Y$  such that for every  $s \in S$   $\pi(y_0 \mid s, s) = 1$  and  $\pi(y_0 \mid s, s') = 0$  for  $s \neq s'$ .

two assumptions on  $\it U$ 

Assumption 2 requires that there is no asymmetric profile that makes both players strictly better off than any symmetric profile.

**Assumption 2:**  $U(a, a') > \max_a U(a, a) \Rightarrow U(a', a) \leq \max_a U(a, a)$ 

with public randomization device Assumption 2 always satisfied if include actions that may depend on the outcome of the public randomization: use coin flip to decide which player is the row player and which player is the column player o

once roles are assigned, players choose the Pareto optimal actions

Assumption 3 requires an action  $\underline{a} \in A$  that ensures that the player gets a payoff that is at least as large as the payoff of his opponent.

**Assumption 3**: There is an  $\underline{a} \in A$  such that  $U(\underline{a}, a) - U(a, \underline{a}) \geq 0$  for all  $a \in A$ .

the payoff difference  $U(a^{\,\prime},a)-U(a,a^{\,\prime})$  defines symmetric zero-sum game hence has a (possibly mixed) minmax strategy

Assumption 3 says the minmax strategy is an element of A: the game defined by the payoff differences has a pure minmax strategy

Assumption 3 always satisfied if we include the possibly mixed minmax action as one of the elements of A

**Benchmark Theorem:** If v = U(a, a) for some  $a \in A$  and  $v \ge u_*$  then v is a Nash equilibrium payoff.

Let  $\overline{a}$  be a Pareto optimal symmetric outcome

$$\overline{a} \in \operatorname{arg\,max}_{a \in A} U(a, a)$$

By assumption 0, there is a strategy  $s_0$  for which.

$$s_0(y) = \begin{cases} \overline{a} & \text{if } y = y_0 \\ \underline{a} & \text{if } y \neq y_0 \end{cases}$$

strategy  $s_0$  takes Pareto efficient action when the opponent uses the same strategy and punishes the opponent by taking action  $\underline{a}$  when the opponent chooses a different strategy

note that punishment action maximizes minimum difference between the player's payoff and his opponent's payoff Theorem 3 shows that long run outcome of the evolutionary dynamics puts positive probability on the strategy  $s_0$ 

every other strategy s that used with positive probability is similar to  $s_0$ : when s meets s both players receive the payoff  $U(\overline{a}, \overline{a})$ ; when s meets  $s_0$  both players receive the same payoff.

**Theorem 3:** Under A0-A3, 
$$\mu(s_0) > 0$$
. If  $\mu(s) > 0$  then  $u(s,s) = u(s_0,s_0) = U(\overline{a},\overline{a})$  and  $u(s,s_0) = u(s_0,s)$ .

if  $\overline{a}$  is the unique symmetric Pareto optimal outcome and if  $U(\underline{a},a)-U(a,\underline{a})>0$  for all  $a\neq\underline{a}$  then  $s_0$  is the unique outcome in the long-run limit

suppose some redundancy description of the game

there may be other strategies that induce the same map as  $s_0$ 

there may be an action  $\widehat{a}$  with  $U(\widehat{a},a)=U(\overline{a},a),$  for all  $a\in A$ 

in either case two or more strategies that satisfy the requirement of  $\boldsymbol{s}_0$ 

strategies differ in extraneous detail only but will not recognize each other as the "same strategy"

long-run distribution places positive weight on every such strategy

during this brief period of transition from one to another players using different versions will punish one another by choosing  $\underline{a}$ , the action that maximizes the difference between the two player's payoff

proof of Theorem 3 shows that the strategy  $s_0$  weakly beats the field that is; weakly beats every other strategy in a pairwise contest  $s_0$  need not be ½ dominant in the ordinary sense underlying game is Prisoner's dilemma  $\tilde{s}$  be a constant strategy that always plays "defect" signals that enable a strategy  $\underline{s}$  to play "defect" against  $s_0$  and "cooperate" against  $\tilde{s}$ 

 $s_{\rm n}$  plays "cooperate" against  $s_{\rm n}$  and "defect" otherwise

against  $\frac{1}{2}s + \frac{1}{2}\underline{s}$  the strategy  $\tilde{s}$  does better than s and therefore s is not  $\frac{1}{2}$  dominant

such a strategy seems to serve no useful purpose except to make  $\overline{s}$  look good against  $s_0$ 

our theory of infrequent innovation provides a rigorous account of why we should not expect such strategies to play a role in determining the long-run equilibrium

they do not themselves do well against  $s_0$  and will not remain around long enough for players to discover that they should play  $\overline{s}$ 

# Gift Exchange and Imperfect Identification

simple additively separable structure

each action a has a cost c(a) and yields a benefit b(a) for the opposing player

payoff of a player who takes action a and whose opponent chooses action a' is

$$U(a,a') = b(a') - c(a)$$

example: two players meet and have an opportunity to exchange goods. c denotes the cost of the good and b describes the benefit of the good for the opposing player

resemble a prisoner's dilemma in that the cost minimizing action is dominant

normalize so that  $c(a) \geq 0$ , with c(a) = 0 for some action  $\hat{a} \in A$  note that this utility function satisfies Assumptions 2 and 3

## Simple Strategies

consider strategies that may not be able to identify with certainty when the opponent is using the same strategy

To keep things simple, we first assume that all strategies use the same symmetric information structure

## **Assumption 4:**

$$\pi(y \mid s, s') = \begin{cases} p(y) & \text{if } s = s' \\ q(y) & \text{if } s \neq s' \end{cases}$$

p(y) probability that a player has type y if he and his opponent use the same strategy

q(y) probability when the two players use different strategies

suppose prior probability a player uses s is  $\alpha$  and that he receives the signal y. posterior probability that the opponent will also play according to s is

$$\frac{\alpha p(y)}{\alpha p(y) + (1 - \alpha)q(y)}.$$

This posterior is greater than  $\alpha$  when p(y)>q(y) and less than  $\alpha$  when q(y)>p(y)

(IC)

**Benchmark Theorem:** The strategy  $s \in S$  is a pure strategy Nash equilibrium if and only if it satisfies

$$\sum_{Y} p(s(y))(b(s(y)) - c(a)) \geq \sum_{Y} q(s(y))b(s(y)), \text{ that is players get more}$$

than the cost minimizing action.

strategy  $s_0$ 

for every signal y the action  $s_0(y)$  solves

$$\max_{a \in A} (p(y) - q(y))b(a) - (p(y) + q(y))c(a)$$

assume the maximization problem has unique solution for every  $\boldsymbol{y}$ 

 $s_0$  rewards opponent when p(y) > q(y)

punishes opponent when q(y) > p(y)

in limiting case where the type allows no inference about play ( p(y)=q(y) ) strategy  $s_0$  minimizes cost c

Theorem 4 shows that in the long run only  $s_0$  is played

**Theorem 4:**  $\mu(s) > 0$  if and only if  $s(y) = s_0(y)$  for all  $y \in Y$ 

signal is y then  $s_0$  maximizes

$$\frac{p(y) - q(y)}{p(y) + q(y)}b(a) - c(a)$$

Let s denote the opponent's strategy choice, and suppose that there a prior of  $\frac{1}{2}$  that the opponent is using  $s_0$ . In that case,

$$\Pr(s = s_0 \mid y) = \frac{p(y)}{p(y) + q(y)}; \Pr(s \neq s_0 \mid y) = \frac{q(y)}{p(y) + q(y)}$$

and therefore

$$\frac{p(y)-q(y)}{p(y)+q(y)} = \Pr(s=s_0 \mid y) - \Pr(s \neq s_0 \mid y)$$

objective function puts a larger weight on the opponent's benefit when he is more likely to use strategy  $s_0$ 

in long-run stable outcome all players are choosing the same strategy probability that opponent using  $s_0$  is one irrespective of signal Nevertheless players punish each other for appearing to be different so equilibrium is typically inefficient

## Complex Strategies

Allow general strategies, but continue to keep strategy  $s_0$ 

consider only strategies  $s_0$  and s being played

strategy  $s_0$  is *informationally superior* to strategy s if the signal generated by  $s_0$  provides better information about the opponent's strategy than the signal generated by s

signal generated by  $s_0$  provides better information (in the sense of Blackwell (1954)) than the signals generated by s if there is a non-negative matrix

$$(\lambda_{yz})_{y\in Y,z\in Y}$$

such that

$$\sum_{y \in Y} \lambda_{yz} = 1, \forall z$$

$$\pi(y \mid s, s) = \sum_{z \in Y} \lambda_{yz} p(z),$$

$$\pi(y \mid s, s_0) = \sum_{z \in Y} \lambda_{yz} q(z);$$

In other words, the signals generated by  $\pi(\cdot \mid s, \cdot)$  are a *garbling* of the signals generated by  $s_0$ 

strategy  $s_0$  is *informationally dominant*, if it is informationally superior to every other strategy s

informational dominance only better information in situations where  $s_0$  and one other competing strategy are played

 $s_0$  may be informationally dominant strategy even though strategy s does better at identifying a third strategy  $\overline{s}$ 

trivial example of an informationally dominant strategy is a strategy that cannot be distinguished from any other strategy

 $\pi(y\mid s,s_0)=\pi(y\mid s,s)$  for all s and hence strategy  $s_0$  is informationally dominant even if strategy  $s_0$  does not generate any information, that is, p(y)=q(y) for all y

strategy  $s_0$  is informationally dominant because it successfully masquerades as other strategies

when strategy  $s_0$  is informationally dominant, it emerges as an outcome of the long-run stable distribution

every strategy that is a long-run stable outcome is similar to strategy  $s_0$ : if  $\mu(s)>0$  then the payoff when s meets s is the same as the payoff when  $s_0$  meets  $s_0$ .

**Theorem 5:** If  $s_0$  informationally dominant then,  $\mu(s_0) > 0$ . Moreover, for every strategies s with  $\mu(s) > 0$  we have  $u(s,s) = u(s_0,s_0)$  and  $u(s,s_0) = u(s_0,s)$ .

we have restricted the informationally dominant strategy to generate symmetric information, that is, generate the same information for every opponent

this allows us to identify a behavior (a map from signals to actions) that is successful against every opponent

if we give up the symmetry assumption we must replace it with a requirement that preserves this ability to do well against all opponents

for example, can assume that there is a reference strategy s such that any signal realization generated by  $s_0$  against an arbitrary opponent is at least as informative as it is against strategy s

Informational dominance would then require that the signal generated against s is informationally superior to the signal generated by any opponent.

Example 1: Perfect identification and additive separability

Against self

$$\max_{a \in A} b(a) - c(a)$$

against difference

$$\max_{a \in A} - b(a) - c(a) = -\min_{a \in A} b(a) + c(a)$$

punishment action minimizes the sum of the player's cost and of the opponent's benefit, so player willing to incur a cost if it leads to a negative payoff to a different opponent

## Example 2.

add the strategy  $\bar{s}$ , which can masquerade as any other strategy and which takes the least cost action  $\hat{a}$  for every signal realization

a player using strategy s cannot determine whether the opponent uses  $\bar{s}$  or s, hence  $\pi(y \mid s, \bar{s}) = \pi(y \mid s, s)$  for all signals  $y \in Y$ .

also players who use  $\overline{s}$  do not receive informative signals about their opponents

such a strategy is informationally dominant

Theorem 5 implies that every strategy that is a long-run stable outcome must play the least cost action

introduction of a cost minimizing strategy that successfully masquerades as other strategies eliminates cooperation between players

Example 3.  $s_0$  need not be  $\frac{1}{2}$  dominant in the ordinary sense

environment of Theorem 5

assume that  $s_0$ , the informationally dominant long-run outcome is not constant

 $\tilde{s}$  be a constant strategy that always plays  $\hat{a}$ 

signals that enable a strategy  $\underline{s}$  to identify  $\tilde{s}$  with certainty and choose an action that maximizes b. Otherwise,  $\underline{s}$  chooses  $\hat{a}$ 

For an appropriate choice of b, the strategy  $\tilde{s}$  does better than s against  $\frac{1}{2}s + \frac{1}{2}\underline{s}$  and therefore  $s_0$  is not  $\frac{1}{2}$  dominant.

*Example 4:* symmetric two-signal scenario,  $Y=\{0,1\}$  and  $p(0)=q(1)=p, p\geq 1/2$ 

signal is y=0 then this is an indication that the two players are using the same strategy

signal is y = 1 it is an indication that the strategies are different

three actions  $a \in \{-1, 0, 1\}$ ,  $b(a) = \beta a, c(a) = 2a^2 + a$ 

trading game with a cooperative action (a = 1), a no-trade action (a = 0), and a hostile action (a = -1)

hostile and the cooperative action are costly for players, whereas the no-trade action is costless

apply Theorem 4 and distinguish the following cases

$$\frac{1}{2p-1} > \beta,$$

then in the unique long run outcome all players to take the no-trade action

$$\beta > \frac{3}{2p-1},$$

then in the unique long run outcome players choose the cooperative action when the signal is  $\ 0$  and the hostile action when the signal is  $\ 1$ 

$$\frac{3}{2p-1} > \beta > \frac{1}{2p-1}$$

then in the unique long run outcome players take the no trade action when the signal is 0 and the hostile action when the signal is 1

in this case, the long run outcome is worse than unique equilibrium of the normal form game

Players choose no trade and hostility and do not realize any of the gains from trade

a significant non-monotonicity

suppose  $9 > \beta > 1$ 

when p = 1 we are in the second, cooperative, case

when p approaches  $\frac{1}{2}$  we are in the first no trade case

but for intermediate values of p, for example p = 5/6, we are in the final case where efficiency is actually less than the autarkic solution that occurs when p approaches  $\frac{1}{2}$ .